


Distal Radius Malunion Treated With Plating and Osteotomy

Case Study

Nathan Edwin Lesley, MD

A 63-year-old patient with a chronic distal radius fracture malunion was treated with the Acumed Acu-Loc 2 Wrist Plating System.




Acumed® is a global leader of innovative orthopaedic and medical solutions.



We are dedicated to developing products, service methods, and approaches that improve patient care.

Case Study | Nathan Edwin Lesley, MD



Patient History

The patient is a 63-year-old female who presented with a chronic malunion of the distal radius from a fracture treated nonoperatively five years earlier. A distal ulna excision had previously been performed for ulnocarpal impaction. The patient reported that she experienced improvement in ulnar-sided symptoms, but continued to complain of wrist pain and loss of wrist flexion. X-rays revealed a malunion with 26 degrees of dorsal tilt and a scapholunate angle of approximately 90 degrees. Preoperative wrist flexion/extension was 20/80. The patient elected to proceed with a distal radius osteotomy.

Treatment

An Acu-Loc 2 plate was provisionally applied distally with K-wires. After confirming proper alignment with fluoroscopy, the plate was applied distally with locking screws. An osteotomy was performed from the radial side using an osteotome. After completion of the osteotomy, the plate was secured to the radial shaft with nonlocking screws. Cancellous chips were placed in the defect.

Postoperative Care

Postoperatively, the patient was placed in a nonremovable wrist splint, followed by a removable wrist splint with gentle ROM instructions beginning at 10 days. At a six-week follow-up, X-rays confirmed osseous union. The patient reported significant improvement from preoperative pain, and her wrist flexion/extension was 40/40. The patient was referred to hand therapy to maximize range of motion.



6 weeks postoperative



HNW70-14-A | Effective: 2018/01 | © 2018 Acumed® LLC

Acumed® Headquarters
5885 NW Cornelius Pass Road
Hillsboro, OR 97124
Office: 888.627.9957
Fax: 503.520.9618
www.acumed.net

These materials contain information about products that may or may not be available in any particular country or may be available under different trademarks in different countries. The products may be approved or cleared by governmental regulatory organizations for sale or use with different indications or restrictions in different countries. Products may not be approved for use in all countries. Nothing contained on these materials should be construed as a promotion or solicitation for any product or for the use of any product in a particular way which is not authorized under the laws and regulations of the country where the reader is located. Specific questions physicians may have about the availability and use of the products described on these materials should be directed to their particular authorized Acumed distributor. Specific questions patients may have about the use of the products described in these materials or the appropriateness for their own conditions should be directed to their own physician.