

## Closed Bimalleolar Ankle Fracture Treatment Using Anatomic Rigid Fixation



Acumed® is a global leader of innovative orthopaedic and medical solutions.

We are dedicated to developing products, service methods, and approaches that improve patient care.



# Case Study | Kent Ellington, MD





### Patient History

The patient is a 53-year-old male who fell five feet off a ladder in July 2015. He sustained a closed bimalleolar fracture that was splinted in the emergency room. He presented to the clinic one day later and went to the operating room post-injury on day three.

#### **Treatment**

Treatment included open reduction internal fixation of his ankle fracture. The Acumed Ankle Plating System 3 and Small Fragment Base Set were used to provide fixation. This required a routine lateral and medial approach to the ankle. The patient was placed in a supine position. The fibula had distal comminution. There was one oblique fracture line that was reduced, then fixed initially with a lag screw. The Lateral Fibula Plate was used to complete ORIF of the fibula. Next, the medial malleolar fragment was reduced, then fixed with two 4.0 mm cannulated partially threaded screws with excellent compression.

### Postoperative Care

At final follow-up, the patient was ambulating well without a brace. He was back to work and walking for exercise.

#### Discussion

This fracture pattern can be difficult to treat due to the fibular comminution. The necessary tools, instruments, plates, and screws required to complete ORIF of ankle fractures are included in the Acumed Ankle Plating System 3. The Lateral Fibula Plate has an excellent fit. With anatomic rigid fixation, the patient was allowed to weight-bear earlier, allowing faster rehabilitation.



10 months postoperative



#### LEX70-06-A | Effective: 2017/05 | © 2017 Acumed® LLC

Acumed® Headquarters 5885 NW Cornelius Pass Road Hillsboro, OR 97124 Office: 888.627.9957 Fax: 503.520.9618 www.acumed.net These materials contain information about products that may or may not be available in any particular country or may be available under different trademarks in different countries. The products may be approved or cleared by governmental regulatory organizations for sale or use with different indications or restrictions in different countries. Products may not be approved for use in all countries. Nothing contained on these materials should be construed as a promotion or solicitation for any product or for the use of any product in a particular way which is not authorized under the laws and regulations of the country where the reader is located. Specific questions physicians may have about the availability and use of the products described on these materials should be directed to their particular authorized Acumed distributor. Specific questions patients may have about the use of the products described in these materials or the appropriateness for their own conditions should be directed to their own physician.