



Cerebral Palsy

Functional Electrical Stimulation

Evidence supporting the efficacy of functional electrical stimulation (FES) to improve gait for children with cerebral palsy (CP) is growing steadily. The results of studies using FES to activate the dorsiflexors during gait are increasingly demonstrating improvements in gait quality, such as gait kinematics and gait symmetry, muscle strength and motor control. Utilizing FES during swing directly impacts the efficacy of gait.¹⁴ Use of peroneal nerve FES to initiate dorsiflexion at terminal stance must be balanced with the child's ability to push off.¹⁵ Studies have contributed results to the evidence demonstrating that push-off is preserved with FES.^{7,15} Based on the outcomes shown in a growing number of studies, peroneal nerve FES may be a valuable treatment option for children with CP.



Benefits of FES found in the published research for children with CP include:

Gait Quality, Speed and Symmetry

- Increased mean dorsiflexion^{13,15}
- Significantly increased peak dorsiflexion^{3-6,9,13,15,16}
- Increased ankle dorsiflexion at initial contact with the use of FES^{4,13}
- Improved heel position at initial contact in some subjects both with and without FES⁵
- Improved ankle and foot posture during stance; more symmetrical heel to toe pattern^{7,11}
- Increased symmetry of step length¹
- Trends toward increased gait velocity and cadence^{1,3,10,19}
- Decreased overall gait deviations compared to normative values for children with CP¹⁰

Patient Preference and Quality of Life

- Children tolerate and/or comply with wearing FES^{11,12,14,15}
- Decreased toe-drag and falls^{14,17}

Strength, Motor Control and Neuroplasticity

- Improved muscle strength and control^{2,9}
- Increased cross sectional area and muscle mass^{2,9}
- Increased voluntary strength^{8,14}
- Improved selective motor control¹⁴
- Decreased spasticity and improved control of the motor patterns used in gait^{8,14}
- Improved Gross Motor Functional Measure scores¹

Energy Cost

- Decreased total work imposed on the ankle in terminal stance^{4,13}
- Decreased energy expenditure¹⁹
- Improved endurance¹⁸

Current evidence for the use of peroneal nerve FES for the treatment of drop foot with children who have CP is very promising in terms of tolerability, kinematic improvements and muscle plasticity. The use of FES with a pediatric population is increasingly becoming a viable and effective standard of care. Technologies like the WalkAide and the Pace XL are at the forefront of this paradigm shift and promise to be an exciting part of future therapeutic interventions.

Bibliography

1. Johnston, T. E., Finson, R. L., McCarthy, J. J., Smith, B. T., Betz, R. R., & Mulcahey, M. J. (2004). Use of Functional Electrical Stimulation to Augment Traditional Orthopaedic Surgery in Children with Cerebral Palsy. *J Pediatr Orthop*, 24, 283–291.
2. Karabay, I., Oztürk, G. T., Malas, F.U., Kara, M., Tiftik, T., Ersöz, M., & Ozçakar, L. (2014). Short-Term Effects of Neuromuscular Electrical Stimulation on Muscle Architecture of the Tibialis Anterior and Gastrocnemius in Children with Cerebral Palsy: Preliminary Results of a Prospective Controlled Study. *Am J Phys Med Rehabil*. [Epub ahead of print]
3. Orlin, M. N., Pierce, S. R., Laughton Stackhouse, C., Smith, B.T., Johnston, T. E., Shewokis, P. A., & McCarthy, J. (2005). Immediate Effect of Percutaneous Intramuscular Stimulation During Gait in Children with Cerebral Palsy: A Feasibility Study. *Dev Med Child Neuro*, 47, 684–690.
4. Pierce, S. R., Laughton, C. A., Smith, B. T., Orlin, M. N., Johnston, T. E., & McCarthy, J. J. (2004). Direct Effect of Percutaneous Electric Stimulation During Gait in Children with Hemiplegic Cerebral Palsy: A Report of 2 Cases. *Arch Phys Med Rehabil*, 85, 339–43.
5. Postans, N. J., & Granat, M. H. (2005). Effect of Functional Electrical Stimulation, Applied During Walking, on Gait in Spastic Cerebral Palsy. *Dev Med Child Neurol*, 47, 46–52.
6. Seifart, A., Unger, M., & Burger, M. (2010). Functional Electrical Stimulation to Lower Limb Muscles After Botox in Children with Cerebral Palsy. *Pediatr Phys Ther*, 22, 199–206.
7. Ho, C. L., Holt, K. G., Saltzman, E., & Wagenaar, R. C. (2006). Functional Electrical Stimulation Changes Dynamic Resources in Children with Spastic Cerebral Palsy. *Phys Ther*, 86, 987–1000.
8. Daichman, J., Johnston, T. E., Evans, K., & Tecklin, J. S. (2003). The Effects of a Neuromuscular Electrical Stimulation Home Program on Impairments and Functional Skills of a Child with Spastic Diplegic Cerebral Palsy: A Case Report. *Pediatr Phys Ther*, 15, 153–158.
9. Damiano, D. L., Prosser, L. A., Curatalo, L.A., & Alter, K. E. (2013). Muscle Plasticity and Ankle Control after Repetitive Use of a Functional Electrical Stimulation Device for Foot Drop in Cerebral Palsy. *Neurorehabil Neural Repair*, 27(3), 200–207.
10. Danino, B., Khamis, S., Hemo, Y., Batt, R., Snir, E., Wientroub, S., & Hayek, S. (2013). The Efficacy of Neuroprosthesis in Young Hemiplegic Patients, Measured by Three Different Gait Indices: Early Results. *J Child Orthop*, 7, 537–542.
11. Durham, S., Eve, L., Stevens, C., & Ewins, D. (2004). Effect of Functional Electrical Stimulation on Asymmetries in Gait of Children with Hemiplegic Cerebral Palsy. *Physiotherapy*, 90(2), 82–90.
12. Meilahn, J. R. (2013). Tolerability and Effectiveness of a Neuroprosthesis for the Treatment of Footdrop in Pediatric Patients with Hemiparetic Cerebral Palsy. *PM R*, 5(6), 503–9.
13. Pierce, S. R., Orlin, M.N., Lauer, R. T., Johnston, T.E., Smith, B.T., & McCarthy, J. J. (2004). Comparison of Percutaneous and Surface Functional Electrical Stimulation During Gait in a Child with Hemiplegic Cerebral Palsy. *Am J Phys Med Rehabil*, 83, 798–805.
14. Pool, D., Blackmore, A. M., Bear, N., & Valentine, J. (2014). Effects of Short-Term Daily Community WalkAide Use on Children with Unilateral Spastic Cerebral Palsy. *Pediatr Phys Ther*, 26(3), 308–17.
15. Prosser, L. A., Curatalo, L. A., Alter, K. E., & Damiano, D.L. (2012). Acceptability and Potential Effectiveness of a Foot Drop Stimulator in Children and Adolescents with Cerebral Palsy. *Dev Med Child Neurol*, 54(11), 1044–1049.
16. van der Linden, M. L., Hazelwood, E., Hillman, S. J., & Robb, J. E. (2008). Functional Electrical Stimulation to the Dorsiflexors and Quadriceps in Children with Cerebral Palsy. *Pediatr Phys Ther*, 20, 23–29.
17. van der Linden, M. (2012). Functional Electrical Stimulation in Children and Adolescents with Cerebral Palsy. *Dev Med Child Neurol*, 54, 972–972. doi:10.1111/j.1469-8749.2012.04419
18. El-Shamy, S. M., & Abdelaal, A. (2016). WalkAide Efficacy on Gait and Energy Expenditure in Children with Hemiplegic Cerebral Palsy: A Randomized Controlled Trial. *AM J Phys Med Rehabil*, 95(9), 629–638.
19. Carroll, M. K., Toelle, C. A., Kim, S. H., Ambler, S. B., & Highsmith, M. J. (2014). The Effect of the WalkAide Functional Electrical Stimulation Unit on Gait Asymmetry in a Child with Cerebral Palsy: A Case Report. *Technology and Innovation*, 15, 287–292.



FES PATIENT SOLUTIONS

4999 Aircenter Circle, Suite 103
 Reno, NV 89502
 888.884.6462 | acplus.com
[Facebook.com/ACPFESPatientSolutions](https://www.facebook.com/ACPFESPatientSolutions)